arXiv:1110.3646v1 [quant-ph] 17 Oct 2011

Density Matrix Recursion Method: Genuine Multisite Entanglement Distinguishes Odd from Even Quantum Heisenberg Ladders

Himadri Shekhar Dhar[1], Aditi Sen(De)[2], and Ujjwal Sen[2]
[1] *School of Physical Sciences, Jawaharlal Nehru University, New Delhi 110 067, India*
[2] *Harish-Chandra Research Institute, Chhatnag Road, Jhunsi, Allahabad 211 019, India*

We introduce an analytical iterative method, the density matrix recursion method, to generate arbitrary reduced density matrices, and bipartite as well as multipartite entanglement of multi-legged periodic spin-1/2 quantum Heisenberg ladders with an arbitrary number of legs. We apply this technique to distinguish between even- and odd-legged ladders. Specifically, we show that while multipartite entanglement decreases with increasing system-size for even-legged ladders, it does the opposite for odd-legged ones.

I. INTRODUCTION

The Heisenberg ladder is an interesting platform to investigate quantum many body systems in the intermediate sector between the one- and two-dimensional lattice structures [1]. The possibility of relating doped even-legged Heisenberg ladders to high-temperature superconductivity [2–4] make such quantum systems extremely important. The characteristic pseudo 2-D structure of ladders evokes considerable interest in several other areas of condensed matter physics [5, 6] and quantum information [7]. On the other hand, concepts like entanglement and quantum correlations have been applied to understand critical quantum phenomena in spin systems [8], including in Heisenberg ladders [7]. Properties of such systems have also been tested experimentally in several systems or proposals thereof have been presented, including in compounds and cold gas [9].

A striking feature of the Heisenberg ladder is that the interpolation from the 1-D spin chain to the 2-D square lattice by gradually increasing the number of legs is not straighforward. The quantum characteristics of the Heisenberg ladder ensures that the odd- and the even-legged ladder ground states have different correlation properties. "Even ladders" have a finite-gapped ground state excitation and exponential decay of two-site correlations, while "odd ladders" are gapless and have power-law decay [6, 10]. The results of Heisenberg ladders cannot therefore be directly extrapolated to the 2-D regime. This difference in quantum characteristic of odd- and even-legged ladders may lead to interesting features in the entanglement properties of the system. However, calculating the bipartite or multipartite entanglement in large-sized multi-legged Heisenberg ladder ground states is numerically challenging. Within a dimer-covering approach [6], it is possible to derive energy density and spin-correlation functions for two-legged ladders by using generating functions [11] or state iterations [12]. Approximate solutions may also be obtained for the four-legged ladder [13].

In this paper, we introduce an analytical iterative technique, the "density matrix recursion method" (DMRM), to obtain the reduced density matrix of an arbitrary number of sites of a quantum spin-1/2 Heisenberg ladder with an arbitrary number of legs and with both open and periodic boundary conditions. Specifically, within a dimer-covering approach, we find separate iterative formalisms for even and odd ladders. These partial density matrices can be used to calculate the scaling and behavior of single-, two-, and multi-site physical properties of the whole system, including two-site correlations and bipartite as well as multipartite entanglement. We then apply our method to obtain the nature of a genuine multipartite entanglement, quantified by the *generalized geometric measure* (GGM) [14]. Multisite entanglement is a useful tool in the topological study of ground states of spin systems [15]. Using a multisite entanglement measure is advantageous over the bipartite correlation measures, as the former involves a greater distribution of information between the multisite sub-regions and the hidden correlations due to tracing out is not negated. We compare odd- and even-legged ladders by using genuine multisite entanglement, and find that the GGM of odd-legged ladders increases with system-size while it decreases in the even ones.

The paper is organized as follows. We introduce the model state in Sec. II. The genuine multiparty entanglement measure is defined in Sec. III. The main results are presented in sections IV and V. In particular, the density matrix recursion methods for even and odd ladders are respectively introduced in sections IV A and IV B. We present a conclusion in Sec. VI.

II. MODEL STATE

Within a dimer-covering approach, the ground state of a antiferromagnetic Heisenberg ladder consists of nearest neighbor dimer pairs on the spin-1/2 lattice. Each bipartite lattice site is occupied by a spin-1/2 qubit. The pairs are limited to nearest neighbor (NN) directed dimer pairs, as governed by the interactions on a *bipartite lattice*. The multi-legged Heisenberg ladder consists of coupled parallel Heisenberg chains with $\mathcal{N}$ sites on each chain labelled from 1 to $\mathcal{N}$. The number of chains, referred as legs, of the ladder are labelled as 1 to M. The total number of spin-1/2 sites in the entire bipartite lattice is s ($s = M\mathcal{N}$). The vertical chains, referred as rungs, are labelled from 1 to n, similar to the sites on the chain, and each rung contains M spins. See Fig. 1. The periodic ladder is conditionally defined by allowing dimer states to be formed between rungs 1 and n [11]. Of course as numbers, $n = \mathcal{N}$. The quantum state under study is therefore [17] $|\mathcal{N}\rangle = \sum_k \left[\prod_{i,j} |(a_i, b_j)\rangle\right]_k$, where $|(a_i, b_j)\rangle$ refers to a dimer between sites a_i and b_j on the bipartite lattice. (i, j) are NN sites with $i \neq j$. A product of all such dimers on the spin-1/2 lattice constitutes a complete dimer covering. The summation refers to the superposition of all such dimer product states or coverings.



FIG. 1. (Color online) The M-legged, $\mathcal{N}$-runged Heisenberg ladder in the form of a *bipartite lattice*. The solid circles are of sublattice A, while the hollow ones are of B. The red arrows (solid) show the nearest-neighbor dimer states ($|(a_i, b_j)\rangle$) from a site in sublattice A to another in B. The arrows (dashed) at the boundary indicate that the lattice is with periodic boundary condition. The figure also shows the two-site ladder state, $|2\rangle$, and the one-site rung state, $|1\rangle$, which will be profusely used in the iteration method presented.

III. GENUINE MULTIPARTITE ENTANGLEMENT

To quantify the amount of genuine multisite entanglement present on multi-legged ladders of different sizes, we consider a genuine multipartite entanglement measure called generalized geometric measure (GGM) [14, 18]. The GGM of an N-party pure quantum state $|\phi_N\rangle$ is defined as

$$\mathcal{E}(|\phi_N\rangle) = 1 - \Lambda^2_{\max}(|\phi_N\rangle),$$

where $\Lambda_{\max}(|\phi_N\rangle) = \max |\langle\chi|\phi_N\rangle|$, with the maximization being over all pure states $|\chi\rangle$ that are not genuinely N-party entangled. An N-party pure quantum state is said to be genuinely multiparty entangled if it is entangled across every bipartite split of the N parties. It was shown in Ref. [14, 18] that $\mathcal{E}(|\phi_N\rangle) = 1 - \max\{\lambda^2_{A:B} | A \cup B = \{1, 2, \ldots, N\}, A \cap B = \emptyset\}$, where $\lambda_{A:B}$ is the maximal Schmidt coefficients in all possible bipartite splits $A : B$ of $|\phi_N\rangle$. The GGM can therefore be calculated once the reduced density matrices of the ladder states can be calculated.

IV. DENSITY MATRIX RECURSION METHOD

The iterative method for generating arbitrary local density matrices are introduced in the following two subsections for the even and the odd ladders. We will show in a subsequent publication that both the iterations developed here can finally lead to obtain an iteration method for the 2D square lattice [19].

A. The Even Ladder

The recursion relation for the non-periodic (i.e. with open boundary conditions) even-legged ladder can be written as

$$\begin{aligned}|\mathcal{N}+2\rangle &= |\mathcal{N}+1\rangle|1\rangle_{n+2} + |\mathcal{N}\rangle|\bar{2}\rangle_{n+1,n+2}\\ &= |\mathcal{N}\rangle|2\rangle_{n+1,n+2} + |\mathcal{N}-1\rangle|\bar{2}\rangle_{n,n+1}|1\rangle_{n+2}, \quad (1)\end{aligned}$$

where $|\mathcal{N}\rangle$ is an even-legged ladder containing $\mathcal{N}$ sites in each chain and corresponds to the rungs numbered from 1 to n. On the other hand, e.g. $|\mathcal{N}\rangle_{2,n+1}$ represents an even-legged ladder containing $\mathcal{N}$ sites in each chain, and corresponds to the rungs numbered from 2 to $n + 1$. $|1\rangle$ is a vertical chain containing M spins representing a single rung of the ladder with the dimer coverings between adjacent sites. $|2\rangle_{n+1,n+2}$ is a two-rung even-legged ladder of rungs $n + 1$ and $n + 2$. Also, $|\bar{2}\rangle_{n+1,n+2} = |2\rangle_{n+1,n+2} - |1\rangle_{n+1}|1\rangle_{n+2}$. If no site numbers are mentioned in subscript of $|\mathcal{N}\rangle$, it is to be considered to be from 1 to $n = \mathcal{N}$.

The periodic boundary condition entails that the ladder also form dimer states between the rungs 1 and n. Accounting for the additional states in the ladder system, due to the periodicity, the recursion relation can be modified into

$$|\mathcal{N}+2\rangle_P = |\mathcal{N}+2\rangle_{1,n+2} + |\mathcal{N}\rangle_{2,n+1}|\bar{2}\rangle_{n+2,1}, \quad (2)$$

where the subscript P stands for a periodic ladder state. Throughout the paper, a state without a subscript P will imply a non-periodic state. The density matrix for the periodic ladder state can be calculated by using (2):

$$\begin{aligned}\rho_P^{(N+2)} &= |\mathcal{N}+2\rangle\langle\mathcal{N}+2|_P\\ &= |\mathcal{N}+2\rangle\langle\mathcal{N}+2| + |\mathcal{N}+2\rangle\langle\mathcal{N}|_{2,n+1}\langle\bar{2}|_{n+2,1}\\ &+ |\mathcal{N}\rangle_{2,n+1}|\bar{2}\rangle_{1,n+2}\langle\mathcal{N}+2|\\ &+ |\mathcal{N}\rangle\langle\mathcal{N}|_{(2,n+1)}|\bar{2}\rangle\langle\bar{2}|_{(1,n+2)} \quad (3)\end{aligned}$$

If, for example, we trace out all but two rungs (say $n+1$ and $n + 2$) from the state, we will obtain a two-rung (mixed state) ladder containing $2M$ spins, where M is the number of legs of the ladder. Note that M is even here.

These reduced states can then be used to calculate the multisite entanglement properties of the ladder system. In particular,

$$\rho^{(2)}_{P(n+1,n+2)} = \mathrm{tr}_{1..n}(|\mathcal{N}+2\rangle\langle\mathcal{N}+2|_P) \tag{4}$$

is the two-rung state of the periodic ladder. Now, using relation (1), we can obtain the trace for the non-periodic part to have $\mathrm{tr}_{1..n}(|\mathcal{N}+2\rangle\langle\mathcal{N}+2|) = \mathrm{tr}_{1..n}[|\mathcal{N}\rangle\langle\mathcal{N}||2\rangle\langle 2|_{(n+1,n+2)} + |\mathcal{N}-1\rangle\langle\mathcal{N}-1||\bar{2}\rangle\langle\bar{2}|_{(n,n+1)}|1\rangle\langle 1|_{(n+2)} + |\mathcal{N}\rangle|2\rangle_{n+1,n+2}\langle\mathcal{N}-1|\langle\bar{2}|_{n,n+1}\langle 1|_{n+2} + |\mathcal{N}-1\rangle|\bar{2}\rangle_{n,n+1}|1\rangle_{n+2}\langle\mathcal{N}|\langle 2|_{n+1,n+2}]$. Tracing over the rungs 1 to n, we get

$$\begin{aligned}\rho^{(2)}_{(n+1,n+2)} &= \mathcal{Z}_{\mathcal{N}}|2\rangle\langle 2|_{(n+1,n+2)} + \mathcal{Z}_{\mathcal{N}-1}\bar{\rho}_{n+1}\otimes|1\rangle\langle 1|_{(n+2)}\\ &+ (|2\rangle_{n+1,n+2}\langle 1|_{n+2}\langle\xi_{\mathcal{N}}|_{n+1} + h.c.)\end{aligned} \tag{5}$$

where $\mathcal{Z}_{\mathcal{N}} = \langle\mathcal{N}|\mathcal{N}\rangle$, $\bar{\rho}_{n+1} = \mathrm{tr}_n[|\bar{2}\rangle\langle\bar{2}|_{(n,n+1)}]$, and $\langle\xi_{\mathcal{N}}|_{n+1} = \langle\bar{2}|_{n,n+1}\langle\mathcal{N}-1|\mathcal{N}\rangle$. Extending the trace to the periodic ladder state, we get

$$\begin{aligned}\rho^{(2)}_{P(n+1,n+2)} &= \rho^{(2)}_{n+1,n+2} + \mathrm{tr}_{1..n}[|\mathcal{N}\rangle\langle\mathcal{N}|_{(2,n+1)}|\bar{2}\rangle\langle\bar{2}|_{(1,n+2)}\\ &\quad + (|\mathcal{N}\rangle_{2,n+1}|\bar{2}\rangle_{1,n+2}\langle\mathcal{N}+2| + h.c.)]\\ &= \rho^{(2)}_{n+1,n+2} + \beta^2_{1(n+1,n+2)} + (\beta^2_{2(n+1,n+2)} + h.c.)\end{aligned} \tag{6}$$

Here, $\beta^2_{1(n+1,n+2)} = \mathcal{Z}_{\mathcal{N}-1}|1\rangle\langle 1|_{(n+1)}\otimes\bar{\rho}_{n+2} + \mathcal{Z}_{\mathcal{N}-2}\bar{\rho}_{n+1}\otimes\bar{\rho}_{n+2} + (|1\rangle\langle\xi_{\mathcal{N}-1}|_{(n+1)}\otimes\bar{\rho}_{n+2} + h.c.)$, with $\langle\xi_{\mathcal{N}-1}|_{n+1} = \langle\bar{2}|_{n,n+1}\langle\mathcal{N}-2|_{2..n-1}\mathcal{N}-1\rangle_{2..n}$. And, $\beta^2_{2(n+1,n+2)} = |2\rangle_{n+1,n+2}\langle 1|_{n+1}\langle\xi_{\mathcal{N}}|_{n+2} + |2\rangle_{n+1,n+2}\langle\phi_{\mathcal{N}}|_{n+1,n+2} + \bar{\rho}_{n+1}\otimes|1\rangle_{n+2}\langle\xi_{\mathcal{N}-1}|_{n+2} + \frac{1}{\mathcal{A}}|\xi_1\rangle_{n+1}|1\rangle_{n+2}\langle 1|_{n+1}\langle\eta_{\mathcal{N}-1}|_{n+2}$, where $\langle\phi_{\mathcal{N}}|_{n+1,n+2} = \langle\bar{2}|_{1,n+2}\langle\bar{2}|_{n,n+1}\langle\mathcal{N}-2|_{2,n-1}\mathcal{N}\rangle$, $\langle\eta_{\mathcal{N}-1}|_{n+2} = \langle\bar{2}|_{1,n+2}\langle\mathcal{N}-1|_{2,n}\mathcal{N}-1\rangle_{1,n-1}|1\rangle_n$, and $\mathcal{A} = \langle 1|1\rangle = \langle\bar{1}|\bar{1}\rangle$.

The local density matrices can be calculated by using the iterations, derived below, of the complete and partial inner products defined above. Using (1), the normalization of the non-periodic ladder is given by

$$\begin{aligned}\mathcal{Z}_{\mathcal{N}} &= \langle\mathcal{N}|\mathcal{N}\rangle\\ &= \mathcal{A}\mathcal{Z}_{\mathcal{N}-1} + \mathcal{B}\mathcal{Z}_{\mathcal{N}-2} + 2\mathcal{C}\mathcal{Y}^1_{\mathcal{N}-2} + 2\mathcal{D}\mathcal{Y}^2_{\mathcal{N}-1}.\end{aligned} \tag{7}$$

where $\bar{\mathcal{A}} = \langle 1|\bar{1}\rangle$ and $\mathcal{B} = \langle\bar{2}|\bar{2}\rangle$. $\mathcal{C}$, $\mathcal{D}$, $\bar{\mathcal{C}}$, and $\bar{\mathcal{D}}$ are given by $\langle 1|\bar{2}\rangle = \mathcal{C}|1\rangle + \mathcal{D}|\bar{1}\rangle$ and $\langle\bar{1}|\bar{2}\rangle = \bar{\mathcal{C}}|1\rangle + \bar{\mathcal{D}}|\bar{1}\rangle$, where $|\bar{1}\rangle$ is a vertical rung with a periodic dimer covering between the topmost and lowermost sites of that rung. The other terms in the normalization can be calculated as follows:

$$\begin{aligned}\mathcal{Y}^1_{\mathcal{N}} &= \langle\mathcal{N}|\mathcal{N}-1\rangle|1\rangle = \mathcal{A}\mathcal{Z}_{\mathcal{N}-1} + \mathcal{C}\mathcal{Y}^1_{\mathcal{N}-1} + \mathcal{D}\mathcal{Y}^2_{\mathcal{N}-1},\\ \mathcal{Y}^2_{\mathcal{N}} &= \langle\mathcal{N}|\mathcal{N}-1\rangle|\bar{1}\rangle = \bar{\mathcal{A}}\mathcal{Z}_{\mathcal{N}-1} + \bar{\mathcal{C}}\mathcal{Y}^1_{\mathcal{N}-1} + \bar{\mathcal{D}}\mathcal{Y}^2_{\mathcal{N}-1}.\end{aligned}$$

The other term in the expression for $\rho^{(2)}_{n+1,n+2}$ that is to be calculated using iterations is

$$\begin{aligned}\langle\xi_{\mathcal{N}}| &= \langle\bar{2}|_{n,n+1}\langle\mathcal{N}-1|\mathcal{N}\rangle\\ &= \langle 1|(\mathcal{C}A^1_{\mathcal{N}} + \bar{\mathcal{C}}A^2_{\mathcal{N}}) + \langle\bar{1}|(\mathcal{D}A^1_{\mathcal{N}} + \bar{\mathcal{D}}A^2_{\mathcal{N}}),\end{aligned} \tag{8}$$

where the iteration variables are given by

$$A^1_{\mathcal{N}} = \sum_{i=1}^{\mathcal{N}}\mathcal{Z}_{\mathcal{N}-i}g_{i-1}, \quad A^2_{\mathcal{N}} = \sum_{i=2}^{\mathcal{N}}\mathcal{Z}_{\mathcal{N}-i}h_{i-1}, \tag{9}$$

with $g_{i+1} = \mathcal{C}g_i + \bar{\mathcal{C}}h_i$, $h_{i+1} = \mathcal{D}g_i + \bar{\mathcal{D}}h_i$, $g_0 = 1$, $h_0 = 0$.

The extra iterative terms in the periodic two-rung density matrix, $\rho^{(2)}_{P(n+1,n+2)}$, can be expressed as

$$\begin{aligned}\langle\phi_{\mathcal{N}}|_{n+1,n+2} &= \langle\bar{2}|_{1,n+2}\langle\bar{2}|_{n,n+1}\langle\mathcal{N}-2|_{2,n-1}\mathcal{N}\rangle\\ &= \langle\bar{2}|_{1,n+2}\langle\bar{2}|_{n,n+1}(\mathcal{X}^{\mathcal{N}}_1|1\rangle_1|1\rangle_n\\ &+ \mathcal{X}^{\mathcal{N}}_2|\bar{1}\rangle_1|1\rangle_n + \mathcal{X}^{\mathcal{N}}_3|1\rangle_1|\bar{1}\rangle_n + \mathcal{X}^{\mathcal{N}}_4|\bar{1}\rangle_1|\bar{1}\rangle_n\\ &+ \langle\bar{2}|_{n-2,n-1}...\langle\bar{2}_{23}|\bar{2}\rangle_{12}...|\bar{2}\rangle_{n-1,n}).\end{aligned} \tag{10}$$

Here, the iterative variables, $\mathcal{X}^N_i$, are defined by using (9), in the following way:

$$\begin{aligned}\mathcal{X}^{\mathcal{N}}_1 &= \sum g_{2i}(A^1_{\mathcal{N}-1-2i} + \mathcal{C}A^1_{\mathcal{N}-2-2i}) + \bar{g}_{2i}\mathcal{D}A^1_{\mathcal{N}-2-2i},\\ \mathcal{X}^{\mathcal{N}}_2 &= \sum g_{2i}(A^2_{\mathcal{N}-1-2i} + \mathcal{C}A^2_{\mathcal{N}-2-2i}) + \bar{g}_{2i}\mathcal{D}A^2_{\mathcal{N}-2-2i},\\ \mathcal{X}^{\mathcal{N}}_3 &= \sum h_{2i}(A^1_{\mathcal{N}-1-2i} + \mathcal{C}A^1_{\mathcal{N}-2-2i}) + \bar{h}_{2i}\mathcal{D}A^1_{\mathcal{N}-2-2i},\\ \mathcal{X}^{\mathcal{N}}_4 &= \sum h_{2i}(A^2_{\mathcal{N}-1-2i} + \mathcal{C}A^2_{\mathcal{N}-2-2i}) + \bar{h}_{2i}\mathcal{D}A^2_{\mathcal{N}-2-2i},\end{aligned} \tag{11}$$

where the summation is from $i = 0$ to $\mathcal{N}$. The variables $\bar{g}_i, \bar{h}_i$ mimics the same relations as g_i, h_i, with initial conditions $\bar{g}_0 = 0$, $\bar{h}_0 = 1$. Similarly, we can derive

$$\begin{aligned}\langle\eta_{\mathcal{N}-1}|_{n+2} &= \langle\bar{2}|_{1,n+2}\langle\mathcal{N}-1|_{2,n}\mathcal{N}-1\rangle_{1,n-1}|1\rangle_n\\ &= (\mathcal{A}A^1_{\mathcal{N}-1} + \bar{\mathcal{A}}A^2_{\mathcal{N}-1})(\mathcal{C}\langle 1| + \mathcal{D}\langle\bar{1}|) + \langle\bar{2}|_{n+2,1}\\ &\quad(\mathcal{X}^{\mathcal{N}-1}_1\langle 1|_2\langle 1|_n + \mathcal{X}^{\mathcal{N}-1}_2\langle\bar{1}|_2\langle 1|_n + \mathcal{X}^{\mathcal{N}-1}_3\\ &\quad\langle 1|_2\langle\bar{1}|_n + \mathcal{X}^{\mathcal{N}-1}_4\langle\bar{1}|_2\langle\bar{1}|_n)|\bar{2}\rangle_{1,2}|1\rangle_n,\end{aligned} \tag{12}$$

where the iterative variables can be derived using (11). These iterations can be used to obtain the partial density matrices of a periodic even-legged ladder, and hence its bipartite as well as genuine multipartite entanglement and other single and multi-site physical quantities, provided the values of $\mathcal{A}$, $\bar{\mathcal{A}}$, $\mathcal{B}$, $\mathcal{C}$, $\bar{\mathcal{C}}$, $\mathcal{D}$, $\bar{\mathcal{D}}$, $\mathcal{Z}_1$, $\mathcal{Y}^1_1$, and $\mathcal{Y}^2_1$ are exactly calculated.

The size of the reduced density matrices depend on the type of the ladder considered. For example, a two-rung reduced density matrix for an M-legged ladder is a $\tau = 2M$-spin matrix. Other reduced density matrices of spins smaller than τ, required e.g. for calculating the GGM, can be obtained from the τ-spin matrix by partial tracings.

B. The Odd Ladder

The periodic recursion for the odd ladder is rather different than the even ladder. This is due to the fact that there exists no analogous state for $|1\rangle$ in the odd ladder. The non-periodic $|\mathcal{N}\rangle$ can be written as a series of $\mathcal{N} = 2$ odd-legged ladders. $|\mathcal{N}\rangle = |2\rangle_{1,2}|2\rangle_{3,4}...|2\rangle_{n-1,n}$. The periodic recursion for the odd M-legged $\mathcal{N}$-spin-per-chain ladder is then given by

$$|\mathcal{N}+2\rangle_P = |\mathcal{N}\rangle_{1,n}|2\rangle_{n+1,n+2} + |\mathcal{N}\rangle_{2,n+1}|2\rangle_{n+2,1} \tag{13}$$

where $|2\rangle_{n+1,n+2}$ is a two-rung odd-legged ladder at rungs $n+1, n+2$. The density matrix can be calculated, as before, using (13): $\rho_P^{N+2} = |\mathcal{N}+2\rangle\langle\mathcal{N}+2|_P$. The reduced density matrices can be obtained by tracing out the requisite number of spins. In particular, for obtaining a two-rung reduced density matrix, we trace out the rungs ranging from 1 to n:

$$\begin{aligned}\rho^{(2)}_{P(n+1,n+2)} &= \mathrm{tr}_{1..n}[|\mathcal{N}+2\rangle\langle\mathcal{N}+2|_P] \\ &= \mathrm{tr}_{1..n}[|\mathcal{N}\rangle\langle\mathcal{N}|_{1,n}|2\rangle\langle 2|_{n+1,n+2} \\ &\quad +|\mathcal{N}\rangle\langle\mathcal{N}|_{2,n+1}|2\rangle\langle 2|_{n+2,1} \\ &\quad +(|\mathcal{N}\rangle_{1,n}|2\rangle_{n+1,n+2}\langle\mathcal{N}|_{2,n+1}\langle 2|_{1,n+2}+h.c.)].\end{aligned}$$

After simplification, the above equation reads

$$\begin{aligned}\rho^{(2)}_{P(n+1,n+2)} &= \mathcal{Z}_{\mathcal{N}}|2\rangle\langle 2|_{(n+1,n+2)} + \mathcal{Z}_{\mathcal{N}-2}\bar{\rho}_{n+1}\otimes\bar{\rho}_{n+2} \\ &\quad + (|2\rangle_{n+1,n+2}\langle\Omega_{\mathcal{N}}|_{n+1,n+2}+h.c.) \qquad (14)\end{aligned}$$

where $\langle\Omega_{\mathcal{N}}|_{n+1,n+2} = \langle 2|_{1,n+2}\langle\mathcal{N}|_{2,n+1}|\mathcal{N}\rangle_{1,n}$, and the normalization is given by

$$\mathcal{Z}_{\mathcal{N}} = \langle\mathcal{N}|\mathcal{N}\rangle = \mathcal{Z}_2^{N/2}. \qquad (15)$$

Now, for a M-legged ladder, $\mathcal{Z}_2$ corresponds to a two-legged M-site-per-chain ladder, and can be calculated from the previous section. The recursion for $\langle\Omega_{\mathcal{N}}|$ is

$$\begin{aligned}\langle\Omega_{\mathcal{N}}|_{n+1,n+2} &= \langle 2|_{1,n+2}\langle\mathcal{N}|_{2,n+1}|\mathcal{N}\rangle_{1,n} \\ &= \langle 2|_{1,n+2}\langle 2|_{2,3}...\langle 2|_{n,n+1}|2\rangle_{1,2}|2\rangle_{3,4}...|2\rangle_{n-1,n}. \quad (16)\end{aligned}$$

Hence the computation involves writing an algorithm to iterate the step $\langle 2|_{i,i+3}\langle 2|_{i+1,i+2}|2\rangle_{i,i+1}$. Once the reduced density matrices are obtained by the iterative method, we can again use them to obtain the different single- and multi-site physical quantities of the system.

V. EVEN VERSUS ODD

To illustrate the effectiveness of the density matrix recursion method, we apply it to obtain a multisite entanglement of multi-legged ladders with both even and odd number of legs. In particular, we consider two- and four-legged ladders among even ladders, and three- and five-legged ladders among odd ones. The iterative variables can be evaluated by using their explicitly calculated initial set of values. The iterations provide the reduced density matrices of the system, which are thereafter utilized to obtain the GGM.

Specifically, for $M=2$, the relevant initial parameters are $\mathcal{Z}_0=1$, $\mathcal{Z}_1=2$, $\mathcal{A}=2$, $\bar{\mathcal{A}}=2$, $\mathcal{C}=1$, $\mathcal{D}=0$, $\bar{\mathcal{C}}=0$, $\bar{\mathcal{D}}=0$, $\mathcal{Y}_1^1=2$, and $\mathcal{Y}_1^2=0$. For $M=4$, the initial parameters are $\mathcal{Z}_0=1$, $\mathcal{Z}_1=4$, $\mathcal{A}=4$, $\bar{\mathcal{A}}=2$, $\mathcal{C}=5$, $\mathcal{D}=1$, $\bar{\mathcal{C}}=2$, $\bar{\mathcal{D}}=3$, $\mathcal{Y}_1^1=4$, and $\mathcal{Y}_1^2=2$. A similar analysis can also be done for higher even-legged ladders.

In a similar fashion, the density matrix recursion method can be applied to obtain the multisite entanglement for odd ladders. The initial parameter required in the recursion for $M=3$ is $\mathcal{Z}_2=44$, and for $M=5$ is $\mathcal{Z}_2=804$.



FIG. 2. (Color online) Genuine multisite entanglement decreases with system-size for even ladders. We perform the iterations for $M=2$ and 4. The iterations are carried out until 18 rungs, i.e. 36 and 72 sites respectively for $M=2$ and 4. However in both the cases, the GGM converges much before those sizes. The vertical axis represents the GGM, while the horizontal one represents the number of rungs. Both axes are dimensionless.



FIG. 3. (Color online) Genuine multisite entanglement increases with system-size for odd ladders. We perform the iterations for $M=3$ and 5. The iterations are until 18 rungs, i.e. 54 and 90 sites respectively for $M=3$ and 5. Again in both the cases, the GGM converges much before 18 rungs. The vertical axis represents the GGM, while the horizontal one represents the number of rungs. Both axes are dimensionless.

We are now ready to compare the multisite entanglements for the even- and odd-legged ladders. The GGMs obtained by the iterative methods clearly capture the characteristic complementary nature of even and odd ladders. We show that the GGM decreases with the increase of system-size in the case of even ladders (Fig. 2). The opposite is true for odd ladders – the GGM increases with system-size (Fig. 3). Although the comparison is made by taking $M=3$ and 5 among odd ladders, we have actually performed the computations also for $M = 1$, which again shows the characteristic increasing GGM of odd ladders. In all the cases, the GGM is calculated by considering reduced density matrices upto $2M$ spins: numerical exact diagonalizations corroborate that considering upto 4 spins is already enough. We have performed the iterative algorithms upto 18 rungs in all the cases (which e.g. is equivalent to 72 spins for the four-legged and 90 spins for the five-legged ladder). As seen in the Figs. 2 and 3, the GGM has already converged much before the maximum number of rungs that we have considered.

VI. CONCLUSION

We have introduced an analytical iterative technique, the density matrix recursion method, which can be efficiently used to obtain arbitrary reduced density matrices of the ground states of spin-1/2 antiferromagnetic quantum Heisenberg ladders with an arbitrary number of legs, within a dimer-covering approach. This technique immediately allows us to obtain single- and multi-site physical properties of the system. In particular, we use the method to obtain the scaling of genuine multisite entanglement in both odd- and even-legged ladders. We find that the genuine multisite entanglement can capture the disparity between even and odd ladders.

ACKNOWLEDGMENTS

The work of HSD is supported by the University Grants Commission (UGC), India under the UGC-Junior Research Fellowship scheme. HSD also thanks the Harish-Chandra Research Institute for hospitality and support during visits.

---

[1] E. Dagatto and T.M. Rice, Science **271**, 618 (1996).

[2] P.W. Anderson, Science **235**, 1196 (1987).

[3] G. Baskaran, Indian J. Phys. **89**, 583 (2006).

[4] M. Greven, R.J. Birgeneau, and U.-J. Wiese, Phys. Rev. Lett. **77**, 1865 (1996).

[5] E. Dagatto, J. Riera, and D. Scalapino, Phys. Rev. B **45**, 5744 (1992)

[6] S.R. White, R.M. Noack, and D.J. Scalapino, Phys. Rev. Lett. **73**, 886 (1994).

[7] J.-L. Song, S.-J. Gu, and H.-Q. Lin, Phys. Rev. B **74**, 155119 (2006); A. B. Kallin, I. Gonzalez, M. B. Hastings, and R. G. Melko, Phys. Rev. Lett. **103**, 117203 (2009); A. Tribedi and I. Bose, Phys. Rev. A **79**, 012331 (2009); I.A. Kovàcs and F. Iglòi, Phys. Rev. B **80**, 214416 (2009); H.S. Dhar and A. Sen(De), arXiv:1003.4401; D. Poilblanc, Phys. Rev. Lett. **105**, 077202 (2010); H. Katsura, N. Kawashima, A.N. Kirillov, V.E. Korepin, S. Tanaka, J. Phys. A: Math. Theor. **43**, 255303 (2010); A.M. Läuchli and J. Schliemann, arXiv:1106.3419, and references therein.

[8] M. Lewenstein, A. Sanpera, V. Ahufinger, B. Damski, A. Sen(De), and U. Sen, Adv. Phys. **56**, 243 (2007); L. Amico, R. Fazio, A. Osterloh, and V. Vedral, Rev. Mod. Phys. **80**, 517 (2008).

[9] S. Ohsugi, K. Magishi, S. Matsumoto, Y. Kitaoka, T. Nagata, and J. Akimitsu Phys. Rev. Lett. **82**, 4715 (1999); J.E. Lorenzo, L.P. Regnault, C. Boullier, N. Martin, S. Vanishri, and C. Marin, Phys. Rev. B **83**, 140413 (2011); A.E. Feiguin and M.P.A. Fisher, *ibid.* **83**, 115104 (2011); M. Dalmonte, P. Zoller, and G. Pupillo, Phys. Rev. Lett. **107**, 163202 (2011), and references therein.

[10] T.M. Rice, S. Gopalan, and M. Sigrist, Europhys. Lett. **23**, 445 (1993).

[11] Y. Fan and M. Ma, Phys Rev B **37**, 1820 (1988).

[12] G. Sierra and M.A. Martin-Delgado, Phys. Rev. B **56**, 8774 (1997).

[13] M. Roncaglia, G. Sierra and M.A. Martin-Delgado, Phys. Rev. B **60**, 12134 (1999).

[14] A. Sen(De) and U. Sen, Phys. Rev. A **81**, 012308 (2010).

[15] A. Kitaev and J. Preskill, Phys. Rev. Lett. **96**, 110404 (2006); M. Levin and X.-G. Wen, Phys. Rev. Lett. **96**, 110405 (2006).

[16] M.M. Wolf, F. Verstraete, M.B. Hastings and J.I. Cirac, Phys. Rev. Lett. **100**, 070502 (2008).

[17] S. Liang, B. Doucot, and P.W. Anderson, Phys. Rev. Lett. **61**, 365 (1988).

[18] A. Sen(De) and U. Sen, arXiv:1002.1253 [quant-ph]

[19] H.S. Dhar, A. Sen(De), and U. Sen, in preparation.